



## Activity-based fate modelling for risk assessment of three ionizable organic compounds (triclosan, furosemide, ciprofloxacin)

Polesel, Fabio; Plósz, Benedek; Trapp, Stefan

*Publication date:*  
2013

*Document Version*  
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

*Citation (APA):*  
Polesel, F., Plósz, B., & Trapp, S. (2013). *Activity-based fate modelling for risk assessment of three ionizable organic compounds (triclosan, furosemide, ciprofloxacin)*. Poster session presented at Micropol & Ecohazard 2013, the 8th IWA Specialist Conference on Assessment and Control of Micropollutants/Hazardous Substances in Water, Zurich, Switzerland.

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



# Activity-based fate modelling for risk assessment of three ionizable organic compounds (triclosan, furosemide, ciprofloxacin)

Fabio Polesel (fabp@env.dtu.dk), Benedek Gy. Plósz, Stefan Trapp

Department of Environmental Engineering, Technical University of Denmark, Bygningstorvet, DK-2800 Kgs. Lyngby, Denmark




## Background and target substances

### Most pharmaceuticals are ionizable compounds

Ionizable chemicals are 49% of all substances registered according to the REACH protocols [1]. 77.5% of pharmaceuticals assessed within a data set of over 500 substances were found to ionize at biologically relevant pH range [2].

### Prediction of the fate of ionizable chemicals: a novel approach


Activity-based models have been developed to predict the environmental fate of organic ionizable compounds, based on their QSAR properties [3,4].





## Objectives and Methods

### Environmental risk assessment of three ionizable pharmaceuticals

- Prediction of fate according to REACH regulation: local ( $C_{local}$ ) and regional concentrations ( $PEC_{regional}$ ) of parent substances
- Combination of steady-state models for WWTP fate (**Activity SimpleTreat**) and regional fate (**MAMI-Multimedia Activity Model for Ionics**) [3,4]
- Assessment of **real emission scenarios** (annual average): Lower Saxony, Southern Sweden, Denmark, Northern Italy
- Validation** of model predictions with literature data [5-12]



## Model predictions



### WWTP fate predictions


- Furosemide had the lowest removal (> 70% discharged via effluent).
- Triclosan was ~50% biodegraded and 25% discharged via effluent.
- Ciprofloxacin was mostly removed in the sludge (>70%) due to very high sorption coefficient.

$C_{local}$  in receiving freshwater bodies were estimated to be  $\geq 0.1 \mu\text{g/L}$  for triclosan and ciprofloxacin. Ciprofloxacin was also found to accumulate in soil ( $C_{local} > 10 \mu\text{g/kg}$ ), as a result of sludge amendment.

$PEC_{regional}$  estimated with MAMI were found to be < 6% of  $C_{local}$  (not shown).

## Risk characterization

$PEC_{local}$  (by combining  $C_{local}$  and  $PEC_{regional}$ ) were used to calculate Risk Characterization Ratios (RCRs). Predicted Non-Effect Concentrations (PNECs) were defined from worst cases in literature or calculated according to REACH TGD. PNECs used were 53 ng/L (triclosan), 3.1  $\mu\text{g/L}$  (furosemide), 5 ng/L (ciprofloxacin) in water and 0.6  $\mu\text{g/kg}$  (triclosan) in soil.



$$RCR = \frac{PEC_{local}}{PNEC}$$

$RCR > 1$  was found for triclosan (1.9–5.1) and ciprofloxacin (2.2–4.3). No potential risk was exhibited to furosemide ( $RCR < 0.2$ ).

## Validation

Results were compared with WWTP removal rates and concentrations in literature to test the reliability of the models.

Substance	WWTP removal			Concentrations in freshwater			
	Simulated	Literature	Ref.	Scenario	$PEC_{local}$	Literature	Ref.
Triclosan	74%–76% (46–48% degraded)	55%–98% (48% degraded)	[5,6]	Lower Saxony	80 ng/L	3–90 ng/L	[9,10]
Furosemide	21%–33%	21%	[7]	Denmark	680 ng/L	250–420 ng/L	[11]
Ciprofloxacin	80%–87% (70%–82% in sludge)	88% (83% in sludge)	[8]	Northern Italy	17–26 ng/L	14–26 ng/L	[12]

## Conclusions

- The modification of the standard fate models (SimpleTreat and Level III regional model) to multispecies models for ionizable compounds was possible and allowed prediction of acids' and zwitterions' fate.
- Model predictions of WWTP fate and of local concentrations were realistic.
- Potentially high risk at local level was associated to triclosan and ciprofloxacin in water, sediment and soil compartments in all scenarios assessed.
- Further investigation is needed, as fate is influenced by e.g., temporal variations of emissions and conjugates' retransformation [13,14].

- References**
- [1] Franco A, Ferranti A, Davidsen C, Trapp S, 2010. *Int J Life Cycle Ass* 15:321–325.
  - [2] Manallack DT 2007. *Perspect Medicin Chem* 1:25–38.
  - [3] Franco A, Song L, Trapp S, 2011. *env.dtu.dk/stt*
  - [4] Franco A, Trapp S, 2010. *Environ Toxicol Chem* 29:789–799.
  - [5] Paxéus N, 2004. *Water Sci Technol* 50:253–260.
  - [6] Heidler J, Halden RU, 2007. *Chemosphere* 66: 362–369.
  - [7] Wahlberg C, Björleinius B, Paxéus N 2011. *Water Sci Technol* 63:1772–1780.
  - [8] Golet EM, Xifra I, Siegrist H, Alder AC, Giger W, 2003. *Environ Sci Technol* 37:3243–3249.
  - [9] Wind T, 2004. *Chemosphere* 54:1145–1153.
  - [10] Bester K, 2005. *Arch Environ Con Tox* 49:9–17.
  - [11] Jacobsen BN, Kjersgaard D, Winther-Nielsen M, Gustavson K, 2004. *Water Sci Technol* 50:37–43.
  - [12] Calamari D, Zuccato E, Castiglioni S, Bagnati R, Fanelli R, 2003. *Environ Sci Technol* 37:1241–1248.
  - [13] Plósz BG, Leknes H, Thomas KV, 2010. *Environ Sci Technol* 44:734–742.
  - [14] Plósz BG, Leknes H, Liltved H, Thomas KV, 2010. *Sci Total Environ* 408:1915–1924.

In all the graphs presented, columns represent the mean value (of % of influent load,  $C_{local}$  and RCR) among all scenarios considered, and error bars refer to the range in which those values are included.

